



AENSI Journals

Australian Journal of Basic and Applied Sciences

Journal home page: www.ajbasweb.com



## Photochemical Purification of Wastewater from the Fungicides and Pesticides Using Advanced Oxidation Processes

<sup>1</sup>Abeer El-Saharty and <sup>2</sup>Ibrahim A. Hassan

<sup>1</sup>National Institute of Oceanography, Academy of Science & Technology, Anfoushi, Alexandria.

<sup>2</sup>Centre of Excellence in Environmental Studies (CEES), King Abdulaziz University, Jeddah, Saudi Arabia.

### ARTICLE INFO

#### Article history:

Received 13 November 2013

Received in revised form 20

December 2013

Accepted 23 December 2013

Available online 25 February 2014

#### Keywords:

Photodegradation, Fungicides

Oxidative process

Fenton reagents, Half-Life, UV

treatment.

### ABSTRACT

Photolysis studies have been carried out in aqueous solutions using UV- photo catalytic system to elucidate the photochemical behavior of some fungicides waters.. HPLC has been used for residues determination. Results clearly exhibited that, carbendazim and metalaxyl were almost stable in the dark while UV irradiation decreased their stabilities. T<sub>1/2</sub> ranged between 2.26 and 8.88 hr. for carbendazim fungicide and from 0.5 to 2.57 hr. for metalaxyl under the conditions of photo-Fenton reactions. The degradation rate of the two tested fungicides were greatly enhanced by Fe(III) than Fe(II). H<sub>2</sub>O<sub>2</sub> accelerated the photodegradation of the two fungicides under photo-Fenton reactions. Consequently, treatment of Fe(III)/ H<sub>2</sub>O<sub>2</sub>/UV gave the highest percentage of dissipation (%D) since, after 360 min of irradiation it reached to 85.50% for carbendazim and 99.98% for metalaxyl. The photodegradation of carbendazim and metalaxyl were greatly enhanced by irradiation under ZnO+H<sub>2</sub>O<sub>2</sub>/UV compared to ZnO/UV alone. The effect of the system ZnO+ H<sub>2</sub>O<sub>2</sub>+UV was as equal as that of the photo-Fenton system FeCl<sub>3</sub>+H<sub>2</sub>O<sub>2</sub>+UV. The half-life values under ZnO/UV were 9.63 and 4.81 hr. for carbendazim and metalaxyl, but the values of T<sub>1/2</sub> were decreased under ZnO+H<sub>2</sub>O<sub>2</sub>/UV to 4.03 and 0.78 hr., respectively. There was a positive relationship between the percentage of fungicides dissipation and the irradiation time for the allrunning treatments.

© 2014 AENSI Publisher All rights reserved.

**To Cite This Article:** Abeer El-Saharty and Ibrahim A. Hassan., Photochemical Purification of wastewater from the Fungicides and pesticides using Advanced Oxidation Processes. *Aust. J. Basic & Appl. Sci.*, 8(1): 434-441, 2014

## INTRODUCTION

Intensive agricultural practices during the last few decades and to the large-scale development of the agrochemical industry caused a dramatic increase in the variety and quantities of agrochemicals present in continental and marine natural waters. Pesticides are one of the most important agrochemicals that contaminant the surface and ground water and their contamination was increased many folds in recent years due to their largescale use in intensive agriculture. The sources of this contamination may be summarized as follows:

(i) pesticide treatment as routine agricultural practice; (ii) rinse water polluted with pesticides from containers and spray equipments; (iii) wastewater from agricultural industry (cleaning or postharvest treatment of fruits and vegetables) and; (iv) plant residues contaminated with pesticides (Devipriya and Yesodharan ,2005).

Advanced oxidation processes (AOPs) are efficient and promising methods to detoxify pesticides, which exhibit chemical stability and resistance to biodegradation (Devipriya and Yesodharan, 2005 and Saien and Khezrianjoo, 2008). The photo-oxidative systems are able to use the UV fraction of sunlight reaching the Earth's surface; however, visible light is not efficiently employed. For this reason, the search for alternative photoactive compounds displaying strong absorption bands in the visible range of the spectrum is still meaningful. (Arques *et al.*, 2009).

Solar-driven processes commonly used in detoxification are the Photo-Fenton system, which is based on a combination of iron salts with hydrogen peroxide (Perez *et al.*, 2006; Evgenidoua *et al.*, 2007; Lapertot *et al.*, 2007 and Martin *et al.*, 2009) and heterogeneous photocatalysis with ZnO (Evgenidoua *et al.*, 2005; Navarro *et al.*, 2009 and Soler *et al.*, 2009). These methods have been shown to be efficient in the elimination of an important number of pesticides (Devipriya and Yesodharan, 2005; Perez *et al.*, 2006; Evgenidoua *et al.*, 2005, 2007; Lapertot *et al.*, 2007; Saien and Khezrianjoo, 2008; Arques *et al.*, 2009, Cernigoj *et al.*, 2010 and Sojic *et al.*, 2010). Carbendazim is a systemic benzimidazole fungicide with an extensive and widespread use.

Analysis of sewage sludge have shown that carbendazim occur in urban water systems (Plagellat, *et al.*, 2004). Under natural environmental conditions, it is very stable and it has been frequently detected in surface

**Corresponding Author:** Abeer El-Saharty, National Institute of Oceanography, Academy of Science & Technology, Anfoushi, Alexandria  
E-mail :saharty@yahoo.com

waters (Fernandez *et al.*, 2001 and Escalada *et al.*, 2006). Meanwhile, it appears as the active substance of benomyl and thiophanate-methyl fungicides and moreover as the main degradation product of these two compounds (Boudina *et al.*, 2003). This product is toxic for humans, animals and plants (Akbarsha *et al.*, 2001). Carbendazim may similarly be found as a pollutant of water resources where it can accumulate. The photodegradation of carbendazim has been studied under different experimental conditions of irradiating with UV light (Panades *et al.*, 2000 and Mazellier *et al.*, 2002a, b and 2003). Metalaxyl is a systemic acylanilide fungicide. It is effective against Oomycetes, specially *Peronospora* such as *Phytophthora* spp., *Pseudoperonospora* spp., *Peronospora* spp., *Sclerospora* spp., *Bremia* spp., *Pythium* spp., and other species causing downy mildews, late blight, damping off as well as root, stem and fruit rots in many crops. It is stable against hydrolysis under normal environmental conditions (Sukul, 2006 and Spessoto *et al.*, 2006). It is also photolytically stable in water and soil when exposed to natural sunlight. Its tolerance to a wide range of pH, light, and temperature leads to its continued use in agriculture (Sukul and Spiteller, 2000). Metalaxyl is considered as a model compound of an organic waste photodegradable substance in water (Topalov *et al.* 1999).

The aim of the present work was to study the effect of different advanced oxidation processes (AOPs), photocatalytic oxidation; Fenton's chemistry (belonging to the class of advanced oxidation processes) and hydrogen peroxide (belonging to the class of chemical oxidation technologies) under UV light irradiation on the degradation of carbendazim and metalaxyl in water solutions to find out the fastest and cheapest degradation system.

## MATERIALS AND METHODS

### Chemicals:

Carbendazim fungicide (C<sub>9</sub>H<sub>9</sub>N<sub>3</sub>O<sub>2</sub>) methyl benzimidazol-2-yl carbamate, [CAS No.:10605-21-7, MW= 191.19] and Metalaxyl (C<sub>15</sub>H<sub>21</sub>NO<sub>4</sub>) methyl *N*-(methoxyacetyl)-*N*-(2,6-xylyl)-DL-alaninate [CAS No: 57837-19-1, MW= 279.3], analytical standard grade 98% and 96% respectively, were supplied by Syngenta Co.

Ferric chloride (FeCl<sub>3</sub>.6H<sub>2</sub>O), Ferrous sulphate (FeSO<sub>4</sub> .7H<sub>2</sub>O), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>30% w/v), zinc oxide (ZnO) and sulphuric acid for pH adjustment were reagents grade. All the used solvents were of A.R. grade.

### Experimental set-up:

For Fenton reagent, the solution was prepared by adding desired amount of carbendazim or metalaxyl (5 mg/L) to water, and then the mixture was carefully agitated in standard flask. Freshly prepared Fe<sup>2+</sup>/H<sub>2</sub>O<sub>2</sub>, Fe<sup>3+</sup>/H<sub>2</sub>O<sub>2</sub> as ferrous sulphate and ferric chloride were then added alone at 20 mg /L or in combination with H<sub>2</sub>O<sub>2</sub> at 0.05%. The initial pH of the solution was adjusted to 2.8, the optimum pH for photo-Fenton treatment using sulphuric acid (Perez *et al.*, 2006 and Pignatello *et al.*, 2006).

For ZnO catalyst, fungicides at concentration of 5 mg/L with the appropriate amount of ZnO catalyst (300 mg/L) was shaken carefully before illumination and left for 30 min in the dark in order to achieve the maximum adsorption of the pesticide onto emiconductors surface, then the pH was adjusted the treatments (Dark, UV light, H<sub>2</sub>O<sub>2</sub>+UV, Fe<sub>2</sub>Cl<sub>3</sub>+UV, Fe<sub>2</sub>Cl<sub>3</sub>+H<sub>2</sub>O<sub>2</sub>+UV, FeSO<sub>4</sub>+UV, FeSO<sub>4</sub>+H<sub>2</sub>O<sub>2</sub>+UV, ZnO+UV and ZnO+H<sub>2</sub>O<sub>2</sub>+ UV) were transferred from the standard flask to a quartz glass cell and exposed to irradiation of the UV mercury lamp (model VL-4.LC '8W' with a wavelength range of 254-365nm, the distance between the lamp and fungicides solution was 15 cm). Samples were taken at different time intervals (10, 30, 60, 120, 240 and 360 min) for HPLC analysis.

### Quantitative estimation of fungicides by HPLC:

The samples were analyzed by HPLC system Perkin Elmer HP 200 L, column C18 reversed A Tracor 985 HPLC and Tracor 951 LC pump. Carbendazim was measured at 285 nm, the used solvent was methanol+H<sub>2</sub>O (4 + 1 by volume) and 10 drops/L ammonium hydroxide. Flow rate was 0-5 ml/min. Metalaxyl was estimated at 205 nm. For the first 15 min the mobile phase (1 ml min<sup>-1</sup>) was 10% acetonitrile in water and pH was adjusted to 4.0 with 0.15M H<sub>3</sub>PO<sub>4</sub>. For the next 5 min., the acetonitrile was increased to 90%. The recovery percentage of carbendazim and metalaxyl was ranged between 95% and 88%, respectively.

### Data Calculation:

The calculation of half-life was performed using the first-order rate equation:

$$C_t = C_0 e^{-kt}$$

Where C<sub>t</sub> represents the concentration at time t; C<sub>0</sub> represents the initial concentration and *k* is the constant rate. When the concentration is reduced to 50% of its initial amount, half-life (T<sub>1/2</sub>) can be determined by the following equation:

$$T_{1/2} = \text{Ln} \frac{C_0}{C} = \left[ \frac{0.897}{k} \right]$$

Where  $k$  is the degradation constant (Monkiedje and Spiteller, 2005).

Where  $C_t$  represents the concentration at time  $t$ ;  $C_0$  represents the initial concentration and  $k$  is the constant rate. When the concentration is reduced to 50% of its initial amount, half-life ( $T_{1/2}$ ) can be determined by the following equation:

$$X = \frac{C_0 - C}{C_0}$$

Where  $C_0$  is the initial concentration of organic compound and  $C$  the concentration at a given time.

#### Statistical Analysis:

Three-way ANOVA was applied using SPSS statistical package (# 4) to test the significance of different treatments. Using light and chemicals as plots and residues as subplots.


## RESULTS AND DISCUSSION

The photodegradation of carbendazim and metalaxyl by UV-irradiation at low wavelength (254 nm) in aqueous solutions using advanced oxidation processes (AOP) by addition of photofenton and photo-catalytic reagent were studied by monitoring their loss by HPLC with UV detection.

#### Degradation of fungicides under dark, UV and hydrogen peroxide/UV light:

The kinetic data derived from first-order function and the half-life of carbendazim and metalaxyl were represented in Tables (1 and 2). The degradation of the fungicides at concentration level of 5 mg/L was observed to estimate their degradation rate under dark, UV light and hydrogen peroxide/UV conditions. In the present study, the absorption spectra of carbendazim and metalaxyl were studied at different times of irradiation (10, 30, 60, 120, 240 and 360 min). The data included the Percentages of dissipation Table (1 and 2) while, the degradation efficiency or conversion of the tested fungicides were presented in (Fig. 1a & b).

The aqueous solution of carbendazim was stable in the dark and nearly no degradation was observed. These results were agreed with the study undertaken by Abu-Qare and Duncan, 2002; Mazellier *et al.*, 2002a and Boudina *et al.*, 2003. The stability of carbendazim was decreased under UV irradiation and the degradation efficiency was 0.186 after 360 min of irradiation (Fig. 2a). In other words, pesticide eliminations could be achieved by using UV irradiation, this was the same result of Shawaqfeh *et al.*, 2010, they found that 50% of Vydine was eliminated under UV (254 nm). The presented data in Table (1) revealed that, the photodegradation and accordingly the percentage of dissipation under H<sub>2</sub>O<sub>2</sub>/UV system was higher than that under UV light alone at the same exposure time, since it reached to 41.87% for H<sub>2</sub>O<sub>2</sub>/UV and 18.56 % for UV alone after 360 min of irradiation. Our finding was agreed with Mazellier *et al.*, 2002a,b and 2003; Chelme-Ayala *et al.*, 2010, who reported that, in the system hydrogen peroxide/UV, the transformation of carbendazim and other pesticides was very efficient because of the generation of hydroxyl radicals in the solution. The observed photoproducts were explained as the result of two main degradation pathways: (1) benzene group oxidation, giving rise to hydroxylated benzimidazoles, and (2) change of the 2-NHCO<sub>2</sub>Me group to 2-NHCO<sub>2</sub>CO<sub>2</sub>H, and 2-NHCO<sub>2</sub>CHO (Mazellier *et al.*, 2002a,b and Boudina *et al.*, 2003). With respect to the significant degradation of carbendazim which was followed by the UV absorbance at appropriate maximum wavelengths, attributed to the benzene ring, and with respect to the reported works on mineralization of other pesticides (Malato *et al.*, 2002; Saien and Khezrianjoo, 2008), the appropriate mineralization reaction can be given here with the most oxidized state as:



**Table 1:** Percentage of dissipation (%D), correlation coefficient (R<sup>2</sup>) and half-life values of carbendazim in aqueous solutions.


Treatment	Percentage of dissipation (%D)						R <sup>2</sup>	T <sub>1/2</sub> (hr.)
	10	20	40	80	160	320		
Dark	0.32	0.32	0.30	0.33	0.34	0.35	0.2359	1570
UV	2.63	8.51	9.19*	11.22*	15.80*	18.56*	0.7301	43.13
H <sub>2</sub> O <sub>2</sub> + UV	3.98	16.87*	21.45*	33.09*	45.98*	61.89*	0.5134	17.83
Fe <sub>2</sub> Cl <sub>3</sub> + UV	11.87*	21.43*	33.63*	39.98	50.09*	70.19*	0.4515	15.98
Fe <sub>2</sub> Cl <sub>3</sub> + H <sub>2</sub> O <sub>2</sub> + UV	15.09*	30.98*	42.01*	52.01*	56.09*	81.01**	0.6728	10.98
ZnO + UV	23.98*	39.09*	47.72*	60.98**	70.98*	89.56**	0.6097	8.92
ZnO + H <sub>2</sub> O <sub>2</sub> + UV	34.97**	42.90**	35.81**	68.76**	82.98***	95.98**	0.5915	3.07

\*=P ≤ 0.05; \*\* = P ≤ 0.01


Slight degradation of metalaxyl was achieved under dark system, the percentage of dissipation was 1.14% after 360 min of irradiation. This degradation might be due to the hydrolysis of this pesticide (Katsumata *et al.*, 2005). Furthermore, approximately 31.24% and 69.34 % of metalaxyl were remediated under UV light and UV/H<sub>2</sub>O<sub>2</sub> system after 360 min, respectively (Table 2). The degradation efficiency was increased using the UV/H<sub>2</sub>O<sub>2</sub> compared with dark system, the values were 0.312 and 0.011, respectively (Fig. 1b). Metalaxyl underwent a parallel/consecutive photodegradation to give *N*-2,6-xylyl-D,L-alaninate (I) and *N*-(methoxyacetyl)-2,6-dimethylaniline (II). Compound (I) was easily converted to 2,6-dimethylaniline (III) as earlier, while compound (II) was more stable to photolysis. (Filippo *et al.*, 1996). The photocatalytic degradation of this fungicide yields its complete mineralization after hydroxylation of the aromatic ring, oxidation of the methylene group and the simultaneous cleavage of the  $\alpha$ -bond from nitrogen to the ester moiety and decarboxylation reactions completed the degradation pathway (Topalov *et al.*, 1999).

In the presence of UV/H<sub>2</sub>O<sub>2</sub>, a first-order rate equation was calculated and the observed half-life were 12.83 and 3.98 hr. for carbendazim and metalaxyl, respectively and these times were shorter than those found in the absence of H<sub>2</sub>O<sub>2</sub> since the values were 23.10 and 16.5 hr., respectively (Table 1 and 2).

The Advanced oxidation process (AOP) is based on the formation of HO• radicals by means of the photolysis of hydrogen peroxide and the subsequent propagation reactions, (Eq. 1) (Marc *et al.*, 2004).



**Fig. 1:** The degradation efficiency of carbendazim (A) and metalaxyl (B) at different irradiation time intervals under dark, UV and H<sub>2</sub>O<sub>2</sub>/UV systems (Each value is a mean of 10 replicates  $\pm$  1 SE).



**Fig. 2:** The degradation efficiency of carbendazim (A) and metalaxyl (B) at different irradiation time intervals under Zinc oxide systems.

**Table 2:** Percentage of dissipation (%D), correlation coefficient (R2) and half-life values of metalaxyl in aqueous solutions.

Treatment	Percentage of dissipation (%D)					R2	T <sub>1/2</sub> (hr.)	
	10	20	40	80	160			320
Dark	0.32	0.32	0.30	0.33	0.34	0.35	0.2359	1570
UV	2.63*	8.51*	9.19*	11.22*	15.80*	18.56*	0.7301	43.13
H <sub>2</sub> O <sub>2</sub> + UV	4.02*	10.11*	14.91	17.89*	20.87	34.98**	0.6728	30.39
Fe <sub>2</sub> Cl <sub>3</sub> + UV	7.09*	12.98*	20.42*	27.01**	33.62	45.02**	0.4358	27.98
Fe <sub>2</sub> Cl <sub>3</sub> + H <sub>2</sub> O <sub>2</sub> + UV	17.87*	24.87	30.71*	40.38**	48.41**	56.34**	0.5672	17.98
ZnO + UV	23.17**	28.98**	35.98**	47.98**	56.78**	67.92**	0.3628	11.98
ZnO + H <sub>2</sub> O <sub>2</sub> + UV	33.98**	41.45**	55.52**	61.92**	70.82**	89.93**	0.8720	6.09

Legends as Table 1.

### Degradation of residues under Photo-Fenton system:

The Fenton method requires H<sub>2</sub>O<sub>2</sub>, Fe<sup>2+</sup> salts and acidic pH. Under these conditions highly reactive and unselective oxidants, OH radicals, are produced. With UV-vis light, the formed Fe<sup>3+</sup> complexes are photolysed, which enables regeneration of Fe<sup>2+</sup> (catalyst): the process is called photo-Fenton (Lapertot *et al.*, 2007).

It was quite clear from Table 1 and 2 that carbendazim and metalaxyl were degraded in shorttime with photo-Fenton reaction, since the T<sub>1/2</sub> ranged between 2.26 and 8.88 hr. for carbendazimfungicide and from 0.5 to 2.57 hr. for metalaxyl. Perusal of these results clearly exhibited that, the degradation rate of the tested fungicides were greatly enhanced by Fe(III) than Fe(II), H<sub>2</sub>O<sub>2</sub> also accelerated the photodegradation of the two fungicides with photo-Fenton reaction. This result was in harmony with the result of Evgenidoua *et al.*, 2007 and Shawaqfeh *et al.*, 2010. Consequently, treatments could be arranged according to the speeding up of the photodegradation of the fungicide carbendazim in decreasing order as follows: Fe(III)/H<sub>2</sub>O<sub>2</sub>/UV > Fe(III)/UV > Fe(II)/H<sub>2</sub>O<sub>2</sub>/UV.

Fe(II)/UV at the different times of irradiation, the percentages of dissipation (%D) of these systems were reached to 85.50, 68.40, 54.77 and 36.87%, respectively after 360 min (Table 1). The values of %D of metalaxyl were higher than those of carbendazim and the treatments followed the same order up to 60 min., while after 120min the order of the treatments was as follow: Fe(III)/H<sub>2</sub>O<sub>2</sub>/UV > Fe(II)/H<sub>2</sub>O<sub>2</sub>/UV > Fe(III)/UV > Fe(II)/UV with values of %D 99.98, 92.92, 88.93 and 81.66%, respectively at 360 min from irradiation (Table 2). There was a positive relationship between the percentage of fungicides dissipation and the irradiation time within the same order of photo-Fenton reagents (Fig. 3a and b). Photo-Fenton had been demonstrated to be more effective than TiO<sub>2</sub> (catalytic agent) (Parra *et al.*, 2000; Bressan *et al.*, 2005 ; Da Hora *et al.*, 2005; Farre *et al.*, 2005; Malato *et al.*, 2005 and Perez *et al.*, 2006).

Fe<sup>2+</sup> salt is rapidly converted to Fe<sup>3+</sup> if there is excess H<sub>2</sub>O<sub>2</sub> (during the first seconds after adding H<sub>2</sub>O<sub>2</sub>), but the •OH produced by Fenton reaction (Eq. 2) are not enough for significant oxidation of organics. When Fe<sup>3+</sup> salt is used, it is reduced to Fe<sup>2+</sup>, mainly by solar irradiation and by H<sub>2</sub>O<sub>2</sub> (Eqs. 3 and 4). The primary step is a ligand-to-metal charge transfer reaction. Later intermediate complexes dissociate as shown in Reaction 5. The ligand can be any Lewis base (L) able to form a complex with ferric iron (OH<sup>-</sup>, H<sub>2</sub>O, HO<sub>2</sub><sup>-</sup>, Cl<sup>-</sup>, R-COO<sup>-</sup>, R-OH, R-NH<sub>2</sub>, *etc.*).

Depending on the reacting ligand, the product may be a hydroxyl radical such as in Eqs. 6 and 7 or another radical derived from the ligand. (Zeppet *et al.*, 1992)



As the system behaves independently of the initial oxidation state of iron, ferrous iron cannot usually be distinguished from ferric iron by the reaction mechanism. However, in practice, differences in degradation rates may be observed (Mazellier *et al.*, 2002b).

### Efficiency of the photocatalyst ZnO:

The changes in carbendazim and metalaxyl concentrations or in other words degradation efficiency at different times during the photocatalytic degradation systems of (ZnO/UV) and (ZnO+H<sub>2</sub>O<sub>2</sub>/UV) were shown in Fig. 2 a and b. Evgenidoua *et al.*, 2005 showed that hydrogen peroxide enhanced the degradation efficiency of the ZnO catalyst and this is due to the dual role of hydrogen peroxide which can act as an electron acceptor promoting the charge separation and it is also able to produce additional •OH radicals. The present results are compatible with them since, the two tested fungicides were more degradable under ZnO+H<sub>2</sub>O<sub>2</sub>/UV system than the other one and metalaxyl had higher values of degradation efficiency than the carbendazim. Also, the positive relationship between the degradation efficiency of the fungicides and the irradiation time was observed.

Tables 1 and 2 showed that, in the presence of ZnO and ZnO/H<sub>2</sub>O<sub>2</sub> under UV light, 35.40 and 83.50% removal of the carbendazim could be achieved in 360 min. While almost the same percentages 44.90 and 84.52% could be reached in the case of fungicide metalaxyl after 60 minutes only under the same conditions.

The first-order rate equation and the half-life ( $T_{1/2}$ ) for carbendazim and metalaxyl were also presented in Tables 1 and 2.  $T_{1/2}$  for carbendazim was 9.63 hours under ZnO/UV system and it was decreased to 2.51 hr. under ZnO+H<sub>2</sub>O<sub>2</sub>/UV system, whilst for metalaxyl it was 4.81 hr. and 0.68 hr. under the two systems, respectively.

The effect of the system ZnO+ H<sub>2</sub>O<sub>2</sub>+UV was equal to the photo-Fenton system FeCl<sub>3</sub>+H<sub>2</sub>O<sub>2</sub>+UV, where the percentages of carbendazim dissipation after 320 min of irradiation were 85.5 and 83.5%, respectively and 99.98 and 99.85% for metalaxyl, respectively.

Irradiation system due to a general ZnO photocatalytic mechanism are proposed as follows:

Under the irradiation of energy greater than the band gap and valence band electrons are promoted to the conduction band leaving a hole behind (Eq. 8). The high oxidative potential of the hole ( $h\nu B^+$ ) in the catalyst permits the direct oxidation of organic matter to reactive intermediates (Eq. 9). Very reactive hydroxyl radicals can also be formed either by the decomposition of water (Eq. 10) or by the reaction of the hole with OH<sup>-</sup> (Eq. 11). The hydroxyl radical is an extremely strong one and a non-selective oxidant that leads to degradation of organic chemicals (Khodja *et al.*, 2001; Daneshvar *et al.*, 2004, 2007 and Behnajady *et al.*, 2006). Electron in the conduction band ( $eCB^-$ ) on the catalyst surface can reduce molecular oxygen to superoxide anion (Eq. 12). This radical, in the presence of organic compound, may form organic peroxides (Eq. 13) or hydrogen peroxide (Eq. 14). Electrons in the conduction band are also responsible for the production of hydroxyl radicals, which have been indicated as the primary cause of organic matter mineralization (Eq. 15) (Daneshvar *et al.*, 2004 and Konstantinou and Albanis, 2004).



The fact that mankind has to face the problem of water pollution as an important threat. So, this study may be used as a step for pesticides elimination from the water and accordingly it can be used to purify wastewater to achieve the desired degree of quality. Moreover, this study may be used as a key step in understanding the interaction between pesticides and foliar fertilization.

## REFERENCES

- Abu-Qare, A.W. and H.J. Duncan, 2002. Photodegradation of the herbicide EPTC and the safener dichlormid, alone and in combination, *Chemosphere*, 46: 1183-1189.
- Akbarsha, M., B. Kadalmani, R. Girija, A. Faridha and K.S. Hamid, 2001. Spermatotoxic effect of carbendazim, *Indian J. Exp. Biol.*, 39: 921-924.
- Arques, A., A.M. Amat, L. Santos-Juanes, R.F. Vercher, M.L. Marin and M.A. Miranda, 2009. Abatement of methidathion and carbaryl from aqueous solutions using organic photocatalysts, *Catalysis Today*, 144: 106-111.
- Behnajady, M.A., N. Modirshahla and Hamzavi, 2006. Kinetic study on photocatalytic degradation of C.I. Acid Yellow 23 by ZnO photocatalyst, *J. Hazard. Mater.*, B133: 226-232.
- benzimidazoles. *Chemosphere*, 65: 237-244.
- Boudina, A., C. Emmelin, A.M.F. Baaliouamer Grenier-Loustalot and J.M. Chovelon, 2003. Photochemical behaviour of carbendazim in aqueous solution, *Chemosphere*, 50: 649-655.
- Bressan, M., L. Liberatorre, N. D'Alessandro, L. Tonucci, C. Belli and G. Ranalli, 2005. *Improved combined chemical and biological treatments of olive oil mill wastewater*, *J. Agric. Food Chem.*, 52: 1228-1233.
- Cernigoj, U., U.L. Stangar and J. Jirkovsky, 2010. Effect of dissolved ozone or ferric ions on photodegradation of thiacloprid in presence of different TiO<sub>2</sub> catalyst, *J. Hazardous Mat.*, 177: 399-406.
- Chelme-Ayala, P., M. Gamal El-Din and D.W. Smith, 2010. Degradation of bromoxynil and trifluralin in natural water by direct photolysis and UV plus H<sub>2</sub>O<sub>2</sub> advanced oxidation process, *Water Res.*, 44: 2221-2228.
- Da Hora, A.E., T.P. Xavier, D.R. de Souza, J.A. de Miranda, E.T. Mendonsa-Duarte, R. Ruggiero, L. de Oliveira and C. Sattler, 2005. Solar photo-Fenton treatment of chip board production waste water. *Sol. Energy*, 77.5: 583-589.

Daneshvar, N., S. Aber, M.S. Seyed Dorraji, A.R. Khataee and M.H. Rasoulifard, 2007. Photocatalytic degradation of the insecticide diazinon in the presence of prepared nanocrystalline ZnO powders under irradiation of UV-C light Rasoulifard Separation and Purification. *Technol.*, 58: 91-98.

Daneshvar, N., D. Salari and A.R. Khataee, 2004. Photocatalytic degradation of azo dye acid red 14 in water on ZnO as an alternative catalyst to TiO<sub>2</sub>, *J. Photochem. Photobiol., A: Chem.*, 162: 317-322.

Devipriya, S. and S. Yesodharan, 2005. Photocatalytic degradation of pesticide contaminants in water, *Solar Energy Mat. & Solar Cells*, 86: 309-348.

Escalada, J.P., A. Pajares, J. Gianotti, W.A. Massad, S. Bertolotti, F. Amat-Guerriand, N.A. Garcia, 2006. Dye-sensitized photodegradation of the fungicide carbendazim and related.

Evgenidou, E., K. Fytianos and I. Poulivos, 2005. Semiconductor-sensitized photodegradation of Dichlorvos in water using TiO<sub>2</sub> and ZnO as catalysts, *App. Catal. B: Environ.*, 59: 81-89.

Evgenidou, E., I. Konstantinou, K. Fytianos and I. Poulivos, 2007. Oxidation of two organophosphorous insecticides by the photo-assisted Fenton reaction, *Water Res.*, 41: 2015-2027.

Farre, M.J., M.I. Franch, S. Malato, J.A. Ayllon, J. Peral, 2005. Domenech. Degradation of some biorecalcitrant pesticides by homogeneous and heterogeneous photocatalytic ozonation, *Chemosphere*, 58: 1127-1133.

Fernandez, M., R. Rodriguez, Y. Pico and J. Manes, 2001. Liquid chromatographic-mass spectrometric determination of post harvest fungicides in citrus fruits, *J. Chromatogr. A*, 912: 301-310.

Filippo, M.P., P. Cabras, V.L. Garau, M. Melis, A. Angioni and D. Bullita, 1996. Photodegradation of pesticides .1. Photolysis rates and half-life of acylanilides and their major metabolites in water, *Toxicol., Environ. Chem.*, 55: 199-214.

Katsumata, H., K. Matsuda, S. Kaneco, T. Suzuki, K. Ohta and Y. Yobiko, 2005. Degradation of carbofuran in aqueous solution by Fe (III) aqua complexes as effective photocatalysts, *J. of photochem. And Photobiol. A: Chem.*, 170: 239-245.

Khodja, A.A., T. Sehili, J. Pilichowski and P. Boule, 2001. Photocatalytic degradation of 2 phenylphenol on TiO<sub>2</sub> and ZnO in aqueous suspension., *Photochem J. Photobiol. A: Chem.*, 141: 231-239.

Konstantinou, I.K. and T.A. Albanis, 2004. TiO<sub>2</sub>-assisted photocatalytic degradation of azo dyes in aqueous solution: kinetic and mechanistic investigation, *Appl. Catal. B: Environ.* 49, pp. 1-14. *J. G. M. Cuppen. P.J. Van den Brink. E. Camps. K. F. Uil. and T.C.M. Brock. Impact of the fungicide carbendazim in fresh water microcosms I. Water quality, breakdown of particulate organic matter and responses of macroinvertebrates, Aquat. Toxicol.*, 48: 233-250.

Lapertot, M., S. Ebrahimi, S. Dazio, A. Rubinelli and D. Pulgarin, 2007. Photo-Fenton and biological integrated process for degradation of a mixture of pesticides, *J. of Photochem. And Photobiol. A: Chemistry*, 186: 34-40.

Malato, S., J. Blanco, J. Caceres, A.R. Fernandez-Alba, A. Aguera and A. Rodriguez, 2002. Photocatalytic treatment of water-soluble pesticides by photo-fenton and TiO<sub>2</sub> using solar energy, *Catal. Today*, 76: 209-220.

Malato, S., J. Blanco, M.I. Maldonado, P. Fernandez, W. Gernjak and I. Oller, 2005. Treatment of chlorinated solvents by TiO<sub>2</sub> photocatalysis and photo-Fenton: influence of operating conditions in a solar pilot plant, *Chemosphere*, 58: 391-398.

Marc, P., G.M. Verónica, A.B. Miguel, G. Jaime and E. Santiago, 2004. Degradation of chlorophenols by means of advanced oxidation processes: a general review, *Appl. Catal. B: Environ.*, 47: 219-256.

Martin, M.M.B., J.A.S. Perez, J.L.G. Sanchez, J.L.C. Lopez and S.M. Rodriguez, 2009. Effect of pesticide concentration on the degradation process by combined solar photo-Fenton and biological treatment, *Water Research*, 43: 3838-3848.

Mazellier, P., E. Leroy and B. Legube, 2002a. Photochemical behavior of the fungicide carbendazim in dilute aqueous solution, *J. Photochem. Photobiol. A Chem.*, 153: 221-227.

Mazellier, P., E. Leroy, J. De Laat and B. Legube, 2003. Degradation of carbendazim by UV/H<sub>2</sub>O<sub>2</sub> investigated by kinetic modeling, *Environ. Chem. Lett.*, 1: 68-72.

Mazellier, P., E. Leroy, J.D. Laat and B. Legube, 2002b. Transformation of carbendazim induced by the H<sub>2</sub>O<sub>2</sub>/UV system in the presence of hydrogenocarbonate ions: involvement of the carbonate radical, *New J. Chem.*, 26: 1784-1790.

Monkiedje, A. and M. Spiteller, 2005. Degradation of metalaxyl and mefenoxam and effects on the microbiological properties of tropical and temperate soils, *Int J Environ Res Public Health*, 2: 272-285.

Navarro, S., J. Fenoll, N. Vela, E. Ruiz and G. Navarro, 2009. Photocatalytic degradation of eight pesticides in leaching water by use of ZnO under natural sunlight, *J. Hazardous Mat.*, 172: 1303-1310.

Panades, R., A. Ibarz and S. Esplugas, 2000. Photodecomposition of carbendazim in aqueous solutions, *Water Res.*, 34: 2951-2954.

Parra, S., V. Sarria, S. Malato, P. Peringer and C. Pulgarin, 2000. Photochemical versus coupled photochemical-biological flow system for the treatment of two biorecalcitrant herbicides: metobromuron and isoproturon, *Appl. Catal. B: Environ.*, 27: 153-168.

Berglof, T., T. Van Dung, H. Kylin and I. Nilsson, 2002. *Carbendazim sorption-desorption in Vietnamese soils*, *Chemosphere*, 48: 267-273.

Perez, M.H., G. Penuela, M.I. Maldonado, O. Malato, P. Fernandez, I. Oller, W. Gernjak and S. Malato, 2006. Degradation of pesticides in water using solar advanced oxidation processes, *Appl. Catal. B: Environ.*, 64: 272-281.

Pignatello, J., E. Oliveros and A. MacKay, 2006. Advanced oxidation processes for organic contaminant destruction based on the Fenton reaction and related chemistry, *Crit. Rev. Environ. Sci. Tech.*, 36: 1-84.

Plagellat, C., T. Kupper, L.F. De Alencastro, D. Grandjean and J. Tarradellas, 2004. Biocides in sewage sludge: quantitative determination in some Swiss wastewater treatment plants, *Bull. Environ. Contam. Toxicol.*, 73: 794-801.

Saien, J. and S. Khezrianjoo, 2008. Degradation of the fungicide carbendazim in aqueous solutions with UV/TiO<sub>2</sub> process: Optimization, kinetics and toxicity studies, *J. Hazardous Mat.*, 157: 269-276.

Shawaqfeh, A.T. and F.A.A. Momani, 2010. Photocatalytic treatment of water soluble pesticide by advanced oxidation technologies using UV light and solar energy, *Solar Energy*. doi: 10.1016/j.solener.2010.03.020.

Sojic Despotovic, D.V., M.I. Abazovic, V.N. Comor, N.D. Abramovic, 2010. Photocatalytic degradation of selected herbicides in aqueous suspensions of doped titania under visible light irradiation, *J. of Hazardous Mat.*, 179: 49-56.

Soler, J., A. Garcia-Ripoll, N. Hayek, P. Miro, R. Vicente, A. Arques and A.M. Amat, 2009. Effect of inorganic ions on the solar detoxification of water polluted with pesticides, *Water Res.*, 43: 4441-4450.

Spessoto, M.A., S.I. Melo and V.L. Ferracini, 2006. Behavior of the fungicide metalaxyl in some Brazilian sandy and clayey soils, *J. of Environ. Sci.*, 20: 262-267.

Sukul, P., 2006. Enzymatic activities and microbial biomass in soil as influenced by metalaxyl residues, *Soil Biology Biochemistry*, 38: 320-326.

Sukul, P. and M. Spiteller, 2000. Metalaxyl persistence, degradation, metabolism, and analytical methods, *Rev Environ Contam Toxicol.*, 164: 1-26.

Topalov, A., D. Molanar-Gabor and J. Csanadi, 1999. Photocatalytic oxidation of the fungicide metalaxyl dissolved in water over TiO<sub>2</sub>, *Water Res.*, 33: 1371-1376.

Zepp, R.G., B.C. Faust and J. Hoigne, 1992. Hydroxyl radical formation in aqueous reactions (pH 3±8) of iron(II) with hydrogen peroxide: the photo-Fenton reaction, *Environ. Sci. Technol.*, 26: 313-319.